

B.Tech 2nd Semester Exam., 2021

(New Course)

MATHEMATICS—II**(Linear Algebra, Transform Calculus and
Numerical Methods)**

Time : 3 hours

Full Marks : 70

Instructions :

- (i) The marks are indicated in the right-hand margin.
(ii) There are **NINE** questions in this paper.
(iii) Attempt **FIVE** questions in all.
(iv) Question No. **1** is compulsory.

1. Answer any seven of the following : $2 \times 7 = 14$

(a) The eigenvalues of a matrix A are 2, 3, 1, then find the eigenvalues of $A^{-1} + A^2$.

(b) If A and B are symmetric matrices, then prove that $AB - BA$ is a skew-symmetric matrix.

(c) Prove that the matrix

$$\frac{1}{3} \begin{bmatrix} -2 & 1 & 2 \\ 2 & 2 & 1 \\ 1 & -2 & 2 \end{bmatrix}$$

is orthogonal.

(d) Prove that $\Delta = \frac{1}{2}\delta^2 + \delta\sqrt{1 + (\delta^2/4)}$.

(e) Find the missing values in the table :

x :	45	50	55	60	65
y :	3	_____	2	_____	-2.4

(f) Use Euler's method to obtain an approximate value of $y(0.4)$ for the equation $y' = x + y$, $y(0) = 1$ with $h = 0.1$.

(g) Obtain the approximate value of $y(1.2)$ for the initial value problem $y' = -2xy^2$, $y(1) = 1$ using Taylor series second-order method with step size $h = 0.1$.

of

$$\frac{s^3}{s^4 - a^4}$$

3/ (a) Verify Cayley-Hamilton theorem for the matrix

$$A = \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$$

Hence compute A^{-1} .

7

(b) Reduce the matrix

$$P = \begin{bmatrix} 1 & 2 & -2 \\ 1 & 2 & 1 \\ -1 & -1 & 0 \end{bmatrix}$$

to diagonal form.

7

4. (a) Find a real root of the equation $x \log_{10} x = 1.2$ by Regula-Falsi method, correct to four decimal places.

7

(b) The following table gives the population of a town during the last six censuses. Estimate the population in 1913 by Newton's forward difference formula :

7

Years	:	1911	1921	1931	1941	1951	1961
Population (in thousands)	:	12	15	20	27	39	52

(i) Evaluate :

$$\int_0^{\infty} e^{-x^2} dx$$

2/ (a) Investigate, for what values of λ and μ do the system of equations $x + y + z = 6$, $x + 2y + 3z = 10$, $x + 2y + \lambda z = \mu$ have (i) no solution, (ii) unique solution and (iii) infinite solutions.

7

(b) Find the rank of the matrix

$$\begin{bmatrix} 3 & 2 & -1 \\ 4 & 2 & 6 \\ 7 & 4 & 5 \end{bmatrix}$$

by reducing it to normal form.

7

(Turn Over)

5/ (a) Find

$$\int_0^6 \frac{e^x}{1+x} dx$$

approximately by using Simpson's 3/8 rule on integration.

7

(b) Evaluate

$$\int_0^8 x \sec x dx$$

using eight intervals by Trapezoidal rule.

7

6. (a) State convolution theorem of Laplace transform and using it find

$$L^{-1} \left\{ \frac{1}{(s^2 + 4)(s + 2)} \right\}$$

7

(b) Use Laplace Transform to solve :

$$\frac{dx}{dt} + y = \sin t, \quad \frac{dy}{dt} + x = \cos t$$

given that $x = 2, y = 0$ at $t = 0$.

7

7. (a) Find the Fourier transform of $f(x)$, defined by

$$f(x) = \begin{cases} 1, & |x| < a \\ 0, & |x| > a \end{cases}$$

and hence evaluate

$$\int_{-\infty}^{\infty} \frac{\sin as \cos sx}{s} ds \quad 10$$

(b) Evaluate :

4

$$\int_0^{\infty} e^{-st} t^3 \sin t dt$$

8. (a) Solve the initial value problem $yy' = x, y(0) = 1$, using the Euler method in $0 \leq x \leq 0.8$, with $h = 0.2$ and $h = 0.1$. Compare the results with the exact solution at $x = 0.8$. Extrapolate the result.

6

(b) Given the initial value problem :

$$\frac{dy}{dx} = \frac{y-x}{y+x}, \quad y(0) = 1$$

Find $y(1)$ by Runge-Kutta fourth-order method taking $h = 0.5$.

8

9. Obtain the approximate value of $y(0.2)$ for the initial value problem $y' = x^2 + y^2$, $y(0) = 1$. Using the methods $y_{n+1} = y_n + hf(x_n, y_n)$, as predictor and

$$y_{n+1} = y_n + \frac{h}{2}[f(x_n, y_n) + f(x_{n+1}, y_{n+1})],$$

as corrector, with $h = 0.1$. Perform two corrector iterations per step.

14
